



Contents lists available at SciVerse ScienceDirect

Journal of Algebra

www.elsevier.com/locate/jalgebra

On the intersection of the \mathcal{F} -maximal subgroups and the generalized \mathcal{F} -hypercentre of a finite group

Wenbin Guo^{a,*}, Alexander N. Skiba^{b,2}^a Department of Mathematics, University of Science and Technology of China, Hefei 230026, PR China^b Department of Mathematics, Francisk Skorina Gomel State University, Gomel 246019, Belarus

ARTICLE INFO

Article history:

Received 9 February 2012

Available online 12 June 2012

Communicated by Gernot Stroth

MSC:

20D10

20D25

Keywords:

 \mathcal{F} -critical group \mathcal{F} -maximal subgroup $\pi\mathcal{F}$ -hypercentre π -Boundary condition

Saturated formation

ABSTRACT

Let \mathcal{F} be a class of groups. A chief factor H/K of a group G is called \mathcal{F} -central in G provided $(H/K) \rtimes (G/C_G(H/K)) \in \mathcal{F}$. We write $Z_{\pi\mathcal{F}}(G)$ to denote the product of all normal subgroups of G whose G -chief factors of order divisible by at least one prime in π are \mathcal{F} -central. We call $Z_{\pi\mathcal{F}}(G)$ the $\pi\mathcal{F}$ -hypercentre of G . A subgroup U of a group G is called \mathcal{F} -maximal in G provided that (a) $U \in \mathcal{F}$, and (b) if $U \leq V \leq G$ and $V \in \mathcal{F}$, then $U = V$. In this paper we study the properties of the intersection of all \mathcal{F} -maximal subgroups of a finite group. In particular, we analyze the condition under which $Z_{\pi\mathcal{F}}(G)$ coincides with the intersection of all \mathcal{F} -maximal subgroups of G .

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

Throughout this paper, all groups are finite and G always denotes a finite group. Moreover p is always supposed to be a prime and π is a non-empty subset of the set \mathbb{P} of all primes. We use \mathcal{G}_π (\mathcal{S}_π) to denote the class of all π -groups (of all soluble π -groups, respectively). In particular, \mathcal{G}_p denotes the class of all p -groups; and we put that $\mathcal{G}_\emptyset = \mathcal{S}_\emptyset = (1)$ is the class of all identity groups. We

* Corresponding author.

E-mail addresses: wbguo@ustc.edu.cn (W. Guo), alexander.skiba49@gmail.com (A.N. Skiba).¹ Research of the first author is supported by a NNSF grant of China (Grant # 11071229) and Wu Wen-Tsun Key Laboratory of Mathematics, USTC, Chinese Academy of Sciences.² Research of the second author is supported by Chinese Academy of Sciences Visiting Professorship for Senior International Scientists (Grant No. 2010T2J12).

also use \mathcal{N} , \mathcal{U} and \mathcal{S} to denote the classes of all nilpotent groups, of all supersoluble groups and of all soluble groups, respectively.

Let \mathcal{F} be a class of groups. A group G is said to be \mathcal{F} -critical if G is not in \mathcal{F} but all proper subgroups of G are in \mathcal{F} [4, p. 517]. If $1 \in \mathcal{F}$, then we write $G^{\mathcal{F}}$ to denote the intersection of all normal subgroups N of G with $G/N \in \mathcal{F}$. For any two classes \mathcal{F} and \mathcal{X} of groups, $\mathcal{X}\mathcal{F}$ is the class of groups G such that $G^{\mathcal{F}} \in \mathcal{X}$.

A formation is a class \mathcal{F} of groups with the following properties: (i) Every homomorphic image of any group in \mathcal{F} belongs to \mathcal{F} ; (ii) If $\mathcal{F} \neq \emptyset$, then $G/G^{\mathcal{F}} \in \mathcal{F}$ for any group G . A formation \mathcal{F} is said to be: *saturated* if $G \in \mathcal{F}$ whenever $G/\Phi(G) \in \mathcal{F}$; *hereditary* if $H \in \mathcal{F}$ whenever $H \leq G \in \mathcal{F}$.

For any formation function $f: \mathbb{P} \rightarrow \{\text{formations of groups}\}$, the symbol $LF(f)$ denotes the collection of all groups G such that either $G = 1$ or $G \neq 1$ and $G/C_G(H/K) \in f(p)$ for every chief factor H/K of G and every $p \in \pi(H/K)$. If $\mathcal{F} = LF(f)$ for some formation function f , then f is said to be a local definition or a *local satellite* (Shemetkov) of \mathcal{F} . Every non-empty saturated formation \mathcal{F} has a unique local satellite F with the following property: For any prime p , both $F(p) \subseteq \mathcal{F}$ and $G \in F(p)$ whenever $G/O_p(G) \in \mathcal{F}$ (see [4, IV, Proposition 3.8]). Such a satellite is called the *canonical local satellite* of \mathcal{F} .

A chief factor H/K of a group G is called \mathcal{F} -central in G provided $(H/K) \rtimes (G/C_G(H/K)) \in \mathcal{F}$. A normal subgroup N of G is said to be $\pi\mathcal{F}$ -hypercentral in G if either $N = 1$ or $N \neq 1$ and every chief factor of G below N of order divisible by at least one prime in π is \mathcal{F} -central in G . The symbol $Z_{\pi\mathcal{F}}(G)$ denotes the $\pi\mathcal{F}$ -hypercentre of G , that is, the product of all normal $\pi\mathcal{F}$ -hypercentral subgroups of G . It is clear that for the \mathcal{F} -hypercentre $Z_{\mathcal{F}}(G)$ of G (see [4, p. 389]) we have $Z_{\mathcal{F}}(G) = Z_{\mathbb{P}\mathcal{F}}(G)$. On the other limited case, when $\pi = \{p\}$, $Z_{p\mathcal{F}}(G)$ is the product of all normal subgroups N of G such that every chief factor of G below N of order divisible by p is \mathcal{F} -central.

A subgroup U of G is called \mathcal{F} -maximal in G provided that (a) $U \in \mathcal{F}$, and (b) if $U \leq V \leq G$ and $V \in \mathcal{F}$, then $U = V$ [4, p. 288]. We denote the intersection of all \mathcal{F} -maximal subgroups of G by $\text{Int}_{\mathcal{F}}(G)$. In the paper [3], Beidleman and Heineken characterized the subgroup $\text{Int}_{\mathcal{F}}(G)$ in the case when G is soluble and \mathcal{F} is a hereditary saturated formation. In this paper, as a development of some results in [12] and [13], we find some new properties and applications of the subgroup $\text{Int}_{\mathcal{F}}(G)$.

Baer [1] proved that $\text{Int}_{\mathcal{N}}(G)$ coincides with the hypercentre $Z_{\infty}(G) = Z_{\mathcal{N}}(G)$ of G . Later, in [9], Sidorov proved that if \mathcal{F} is the class of all soluble groups G of nilpotent length $l(G) \leq r$ ($r \in \mathbb{N}$), then for each soluble group G , the equality $Z_{\mathcal{F}}(G) = \text{Int}_{\mathcal{F}}(G)$ holds. In the papers [12] and [13], the analogous results were obtained for the classes of all p -decomposable groups and of all groups G with $G' \leq F(G)$ in the universe of all groups. As one of our results in this paper, we shall also prove that the intersection of all maximal p -nilpotent subgroups of G coincides with the subgroup $Z_{p\mathcal{N}}(G)$. But in general, $Z_{\pi\mathcal{F}}(G) \neq \text{Int}_{\mathcal{F}}(G)$, even when \mathcal{F} is the class of all supersoluble (all p -supersoluble, for any odd prime p) groups and G is soluble (see Theorem A and Remark 4.8 in Section 4).

Definition 1.1. Let \mathcal{X} be a non-empty class of groups and $\mathcal{F} = LF(F)$ be a saturated formation, where F is the canonical local satellite of \mathcal{F} . We say that \mathcal{F} satisfies the π -boundary condition (the boundary condition if $\pi = \mathbb{P}$) in \mathcal{X} if $G \in \mathcal{F}$ whenever $G \in \mathcal{X}$ and G is an $F(p)$ -critical group for at least one $p \in \pi$.

We say that \mathcal{F} satisfies the π -boundary condition if \mathcal{F} satisfies the π -boundary condition in the class of all groups.

If \mathcal{F} is a non-empty formation with $\pi(\mathcal{F}) = \emptyset$, then $\mathcal{F} = (1)$, and therefore for any group G we have $Z_{\mathcal{F}}(G) = 1 = \text{Int}_{\mathcal{F}}(G)$. In the other limited case, when $\mathcal{F} = \mathcal{G}$ is the class of all groups, we have $Z_{\mathcal{F}}(G) = G = \text{Int}_{\mathcal{F}}(G)$. Similarly, if $\mathcal{F} = \mathcal{S}$, then $Z_{\mathcal{F}}(G) = G = \text{Int}_{\mathcal{F}}(G)$ for every soluble group G .

For the general case, we shall prove the following.

Theorem A. Let \mathcal{F} be a hereditary saturated formation with $(1) \neq \mathcal{F} \neq \mathcal{G}$. Let $\pi \subseteq \pi(\mathcal{F})$. Then the equality

$$Z_{\pi\mathcal{F}}(G) = \text{Int}_{\mathcal{F}}(G)$$

holds for each group G if and only if $\mathcal{N} \subseteq \mathcal{F} = \mathcal{G}_{\pi'}\mathcal{F}$ and \mathcal{F} satisfies the π -boundary condition.

Note that $N(p) = \mathcal{G}_p$, where N is the canonical local satellite of \mathcal{N} . Hence every $N(p)$ -critical group has prime order. This shows that \mathcal{N} satisfies the boundary condition and so the above-mentioned Baer's result is a corollary of Theorem A.

Theorem B. Let \mathcal{F} be a hereditary saturated formation of soluble groups with $(1) \neq \mathcal{F} \neq \mathcal{S}$. Let $\pi \subseteq \pi(\mathcal{F})$. Then the equality

$$Z_{\pi\mathcal{F}}(G) = \text{Int}_{\mathcal{F}}(G)$$

holds for each soluble group G if and only if $\mathcal{N} \subseteq \mathcal{F} = \mathcal{S}_{\pi}\mathcal{F}$ and \mathcal{F} satisfies the π -boundary condition in the class of all soluble groups.

If for some classes \mathcal{F} and \mathcal{M} of groups we have $\mathcal{F} \subseteq \mathcal{M}$, then every \mathcal{F} -maximal subgroup of a group is contained in some its \mathcal{M} -maximal subgroup. Nevertheless, the following example shows that in general, $\text{Int}_{\mathcal{F}}(G) \not\subseteq \text{Int}_{\mathcal{M}}(G)$.

Example 1.2. Let $\mathcal{F} = \mathcal{U}$ and \mathcal{M} be the class of all p -supersoluble groups, where $p > 2$. Let q be a prime dividing $p - 1$ and $G = P \rtimes (Q \rtimes C)$, where C is a group of order p , Q is a simple $\mathbb{F}_q G$ -module which is faithful for C and P is a simple $\mathbb{F}_p G$ -module which is faithful for $Q \rtimes C$. Then, clearly, $P = \text{Int}_{\mathcal{F}}(G)$ and $\text{Int}_{\mathcal{M}}(G) = 1$.

This example is a motivation for the following our result.

Theorem C. Let $\mathcal{F} \subseteq \mathcal{M} = LF(M)$ be hereditary saturated formations with $\pi \subseteq \pi(\mathcal{F})$, where M is the canonical local satellites of \mathcal{M} .

(a) Suppose that $\mathcal{N} \subseteq \mathcal{M} = \mathcal{G}_{\pi}\mathcal{M}$ and \mathcal{F} satisfies the π -boundary condition in \mathcal{M} . Then the inclusion

$$\text{Int}_{\mathcal{F}}(G) \leq \text{Int}_{\mathcal{M}}(G)$$

holds for each group G .

(b) If every (soluble) $M(p)$ -critical group belongs to \mathcal{F} for every $p \in \pi$, then $\mathcal{N} \subseteq \mathcal{M}$ and

$$\text{Int}_{\mathcal{F}}(G) \leq Z_{\pi\mathcal{M}}(G)$$

for every (soluble) group G .

Recall that a subgroup H of a group G is said to be \mathcal{F} -subnormal (in the sense of Kegel [8]) or K - \mathcal{F} -subnormal in G (see p. 236 in [2]) if either $H = G$ or there exists a chain of subgroups

$$H = H_0 < H_1 < \cdots < H_t = G$$

such that either H_{i-1} is normal in H_i or $H_i/(H_{i-1})_{H_i} \in \mathcal{F}$ for all $i = 1, \dots, t$.

For any group G , we write $\text{Int}_{\mathcal{F}}^*(G)$ to denote the intersection of all non- K - \mathcal{F} -subnormal \mathcal{F} -maximal subgroups of G . The following theorem shows that for any hereditary saturated formation \mathcal{F} with $\mathcal{N} \subseteq \mathcal{F}$, the intersection of all non- K - \mathcal{F} -subnormal \mathcal{F} -maximal subgroups of a group G coincides with $\text{Int}_{\mathcal{F}}(G)$.

Theorem D. Let \mathcal{F} be a hereditary saturated formation containing all nilpotent groups. Then the equality

$$\text{Int}_{\mathcal{F}}^*(G) = \text{Int}_{\mathcal{F}}(G)$$

holds for each group G .

We prove Theorems A, B, C and D in Section 3. In Section 4 we give some examples and discuss applications of these theorems.

All unexplained notation and terminology are standard. The reader is referred to [4,2,6] if necessary.

2. Preliminaries

In view of Proposition 3.16 in [4, IV], we have

Lemma 2.1. *Let $\mathcal{F} = LF(F)$ be a hereditary saturated formation, where F is the canonical local satellite of \mathcal{F} . Then for any prime p , the formation $F(p)$ is hereditary.*

We shall need some facts about the $\pi\mathcal{F}$ -hypercentre in our proofs.

Lemma 2.2. *Let $\mathcal{F} = LF(F)$ be a saturated formation, where F is the canonical local satellite of \mathcal{F} . Let $\pi \subseteq \pi(\mathcal{F})$ and $\sigma = \pi(\mathcal{F}) \setminus \pi$. Let N be a normal subgroup of G , and $A \leq G$.*

- (1) *A chief factor H/K of G is \mathcal{F} -central if and only if $G/C_G(H/K) \in F(p)$ for all primes p dividing $|H/K|$.*
- (2) *Every G -chief factor of $Z_{\pi\mathcal{F}}(G)$ of order divisible by at least one prime in π is \mathcal{F} -central.*
- (3) $Z_{\pi\mathcal{F}}(G)N/N \leq Z_{\pi\mathcal{F}}(G/N)$.
- (4) $Z_{\pi\mathcal{F}}(A)N/N \leq Z_{\pi\mathcal{F}}(AN/N)$.
- (5) *If \mathcal{F} is (normally) hereditary and H is a (normal) subgroup of G , then $Z_{\pi\mathcal{F}}(H) \cap A \leq Z_{\pi\mathcal{F}}(H \cap A)$.*
- (6) *If $\mathcal{G}_\sigma\mathcal{F} = \mathcal{F}$ and $G/Z_{\pi\mathcal{F}}(G) \in \mathcal{F}$, then $G \in \mathcal{F}$.*
- (7) *Suppose that \mathcal{F} is (normally) hereditary and let H be a (normal) subgroup of G . If $\mathcal{G}_\sigma\mathcal{F} = \mathcal{F}$ and $H \in \mathcal{F}$, then $Z_{\pi\mathcal{F}}(G)H \in \mathcal{F}$.*

Proof. (1) This assertion is well known (see for example Theorem 17.14 in [11] or Theorem 3.1.6 in [6]). Assertions (2) and (6) are evident.

(3) Let H/K be a chief factor of G such that $N \leq K < H \leq NZ_{\pi\mathcal{F}}(G)$ and $|H/K|$ is divisible by at least one prime in π . Then H/K is G -isomorphic to the chief factor $H \cap Z_{\pi\mathcal{F}}(G)/K \cap Z_{\pi\mathcal{F}}(G)$ of G . Therefore H/K is \mathcal{F} -central in G by (1) and (2). Consequently, $Z_{\pi\mathcal{F}}(G)N/N \leq Z_{\pi\mathcal{F}}(G/N)$.

(4) Let $f: A/A \cap N \rightarrow AN/N$ be the canonical isomorphism from $A/A \cap N$ onto AN/N . Then $f(Z_{\pi\mathcal{F}}(A/A \cap N)) = Z_{\pi\mathcal{F}}(AN/N)$ and

$$f(Z_{\pi\mathcal{F}}(A)(A \cap N)/(A \cap N)) = Z_{\pi\mathcal{F}}(A)N/N.$$

By (3) we have

$$Z_{\pi\mathcal{F}}(A)(A \cap N)/(A \cap N) \leq Z_{\pi\mathcal{F}}(A/A \cap N).$$

Hence $Z_{\pi\mathcal{F}}(A)N/N \leq Z_{\pi\mathcal{F}}(AN/N)$.

(5) First suppose that \mathcal{F} is hereditary. Let

$$1 = Z_0 < Z_1 < \cdots < Z_t = Z_{\pi\mathcal{F}}(G)$$

be a chief series of G below $Z_{\pi\mathcal{F}}(G)$ and $C_i = C_G(Z_i/Z_{i-1})$. Let q be a prime divisor of

$$|Z_i \cap H/Z_{i-1} \cap H| = |Z_{i-1}(Z_i \cap H)/Z_{i-1}|.$$

Suppose that q divides $|Z_i \cap H/Z_{i-1} \cap H|$. Then q divides $|Z_i/Z_{i-1}|$, so $G/C_i \in F(q)$ by (1). Hence $H/H \cap C_i \simeq C_i H/C_i \in F(q)$. But $H \cap C_i \leq C_H(Z_i \cap H/Z_{i-1} \cap H)$. Hence $H/C_H(Z_i \cap H/Z_{i-1} \cap H) \in F(q)$ for all primes q dividing $|Z_i \cap H/Z_{i-1} \cap H|$. Thus $Z_{\pi\mathcal{F}}(G) \cap H \leq Z_{\pi\mathcal{F}}(H)$ by (1) and (2). But then

$$Z_{\pi\mathcal{F}}(H) \cap E = Z_{\pi\mathcal{F}}(H) \cap (H \cap E) \leq Z_{\pi\mathcal{F}}(H \cap E).$$

Similarly, one may prove the second assertion of (5).

(7) Since $H \in \mathcal{F}$ we have

$$HZ_{\pi\mathcal{F}}(G)/Z_{\pi\mathcal{F}}(G) \simeq H/H \cap Z_{\pi\mathcal{F}}(G) \in \mathcal{F}$$

and

$$Z_{\pi\mathcal{F}}(G) \leq Z_{\pi\mathcal{F}}(Z_{\pi\mathcal{F}}(G)H)$$

by (5). Hence $HZ_{\pi\mathcal{F}}(G) \in \mathcal{F}$ by (6).

The lemma is proved. \square

The following lemma is evident (note only that statement (i) directly follows from [4, Theorem A.9.2(c)]).

Lemma 2.3. Let \mathcal{F} be a hereditary saturated formation. Let $N \leq U \leq G$, where N is a normal subgroup of G .

- (i) If $G/N \in \mathcal{F}$ and V is a minimal supplement of N in G , then $V \in \mathcal{F}$.
- (ii) If U/N is an \mathcal{F} -maximal subgroup of G/N , then $U = U_0N$ for some \mathcal{F} -maximal subgroup U_0 of G .
- (iii) If V is an \mathcal{F} -maximal subgroup of U , then $V = H \cap U$ for some \mathcal{F} -maximal subgroup H of G .

The proofs of our theorems are based on the following general facts on the subgroup $\text{Int}_{\mathcal{F}}(G)$.

Lemma 2.4. Let \mathcal{F} be a hereditary saturated formation, $\pi \subseteq \pi(\mathcal{F})$ and $\sigma = \pi(\mathcal{F}) \setminus \pi$. Let H, E be subgroups of G , N a normal subgroup of G and $I = \text{Int}_{\mathcal{F}}(G)$. Then:

- (a) $\text{Int}_{\mathcal{F}}(H)N/N \leq \text{Int}_{\mathcal{F}}(HN/N)$.
- (b) $\text{Int}_{\mathcal{F}}(H) \cap E \leq \text{Int}_{\mathcal{F}}(H \cap E)$.
- (c) If $H/H \cap I \in \mathcal{F}$, then $H \in \mathcal{F}$.
- (d) If $H \in \mathcal{F}$, then $IH \in \mathcal{F}$.
- (e) If $N \leq I$, then $I/N = \text{Int}_{\mathcal{F}}(G/N)$.
- (f) $\text{Int}_{\mathcal{F}}(G/I) = 1$.
- (g) If $\mathcal{G}_{\sigma}\mathcal{F} = \mathcal{F}$, then $Z_{\pi\mathcal{F}}(G) \leq I$.

Proof. Assertions (a)–(f) are proved in [12]. Now we prove (g). Let H be a subgroup of G such that $H \in \mathcal{F}$. Then $HZ_{\pi\mathcal{F}}(G)/Z_{\pi\mathcal{F}}(G) \simeq H/H \cap Z_{\pi\mathcal{F}}(G) \in \mathcal{F}$ and $Z_{\pi\mathcal{F}}(G) \leq Z_{\pi\mathcal{F}}(HZ_{\pi\mathcal{F}}(G))$ by Lemma 2.2(5). It follows from Lemma 2.2(5) that $HZ_{\pi\mathcal{F}}(G) \in \mathcal{F}$. Thus $Z_{\pi\mathcal{F}}(G) \leq I$. \square

The following lemma is a corollary of general results on f -hypercentral action (see [4, Chapter IV, Section 6]). For the reader's convenience, we give a direct proof.

Lemma 2.5. Let $\mathcal{F} = LF(F)$ be a saturated formation, where F is the canonical local satellite of \mathcal{F} . Let E be a normal p -subgroup of G . If $E \leq Z_{\mathcal{F}}(G)$, then $G/C_G(E) \in F(p)$.

Proof. Let $1 = E_0 < E_1 < \dots < E_t = E$ be a chief series of G below E . Let $C_i = C_G(E_i/E_{i-1})$ and $C = C_1 \cap \dots \cap C_t$. Then $C_G(E) \leq C$ and so $C/C_G(E)$ is a p -group by Corollary 3.3 in [5, Chapter 5]. On the other hand, by Lemma 2.2(1), $G/C_i \in F(p)$, so $G/C \in F(p)$. Hence $G/C_G(E) \in F(p) = \mathcal{G}_p F(p)$. The lemma is proved. \square

Lemma 2.6. Let $\mathcal{F} = LF(F)$ and \mathcal{M} be saturated formations with $p \in \pi(\mathcal{F})$ and $\mathcal{F} \subseteq \mathcal{M}$, where F is the canonical local satellite of \mathcal{F} . Suppose that G is a group of minimal order in the set of all $F(p)$ -critical groups $G \in \mathcal{M}$ with $G \notin \mathcal{F}$. Then $O_p(G) = 1 = \Phi(G)$ and $G^{\mathcal{F}}$ is the unique minimal normal subgroup of G .

Proof. Let N be a minimal normal subgroup of G . First we show that $G/N \in \mathcal{F} \cap \mathcal{M}$. Indeed, since $G \in \mathcal{M}$ and \mathcal{M} is a formation, $G/N \in \mathcal{M}$. Suppose that $G/N \notin \mathcal{F}$. Then $G/N \notin F(p)$ since $F(p) \subseteq \mathcal{F}$.

On the other hand, for any maximal subgroup M/N of G/N we have $M/N \in F(p)$ since $F(p)$ is a formation and G is an $F(p)$ -critical group. Thus G/N is an $F(p)$ -critical group with $G/N \notin \mathcal{M} \setminus \mathcal{F}$, which contradicts the minimality of G . Hence $G/N \in \mathcal{F}$. Since \mathcal{F} is a saturated formation, $N = G^{\mathcal{F}}$ is a unique minimal normal subgroup of G and $\Phi(G) = 1$. Suppose that $N \leq O_p(G)$ and let M be a maximal subgroup of G such that $G = NM$. Then $G/N \simeq M/N \cap M \in F(p) = \mathcal{S}_p F(p)$, and so $G \leq F(p) \subseteq \mathcal{F}$. This contradiction shows that $O_p(G) = 1$. The lemma is proved. \square

Lemma 2.7. *Let \mathcal{F} be a formation, H and E be subgroups of a group G , where H is K - \mathcal{F} -subnormal in G . Then:*

- (i) $H \cap E$ is K - \mathcal{F} -subnormal in E (see Theorem 6.1.7 in [2]).
- (ii) If E is normal in G , then HE/E is K - \mathcal{F} -subnormal in G/E (see Theorem 6.1.6 in [2]).

Lemma 2.8. *Let \mathcal{F} be a hereditary saturated formation. Let $N \leq U \leq G$, where N is a normal subgroup of G .*

- (i) *If U/N is a non- K - \mathcal{F} -subnormal \mathcal{F} -maximal subgroup of G/N , then $U = U_0 N$ for some non- K - \mathcal{F} -subnormal \mathcal{F} -maximal subgroup U_0 of G .*
- (ii) *If V is a non- K - \mathcal{F} -subnormal \mathcal{F} -maximal subgroup of U , then $V = H \cap U$ for some non- K - \mathcal{F} -subnormal \mathcal{F} -maximal subgroup H of G .*

Proof. (i) By Lemma 2.3(ii), there is an \mathcal{F} -maximal subgroup U_0 of G such that $U = U_0 N$. Since U/N is non- K - \mathcal{F} -subnormal in G/N , U_0 is not K - \mathcal{F} -subnormal in G by Lemma 2.7(ii).

(ii) By Lemma 2.3(iii), for some \mathcal{F} -maximal subgroup H of G we have $V = H \cap U$. Since V is a non- K - \mathcal{F} -subnormal in U , H is not K - \mathcal{F} -subnormal in G by Lemma 2.7(i). \square

Lemma 2.9. *Let $\mathcal{F} = LF(F)$ be a non-empty saturated formation, where F is the canonical local satellite of \mathcal{F} .*

- (1) *If $\mathcal{F} = \mathcal{S}_p \mathcal{F}$ for some prime p , then $F(p) = \mathcal{F}$.*
- (2) *If $\mathcal{F} = \mathcal{NH}$ for some non-empty formation \mathcal{H} , then $F(p) = \mathcal{S}_p \mathcal{H}$ for all primes p .*

Proof. (1) Since $F(p) \subseteq \mathcal{F}$, we need only prove that $\mathcal{F} \subseteq F(p)$. Suppose that this is false and let A be a group of minimal order in $\mathcal{F} \setminus F(p)$. Then $A^{F(p)}$ is a unique minimal normal subgroup of A since $F(p)$ is a formation. Moreover, $O_p(A) = 1$ since $F(p) = \mathcal{S}_p F(p)$. Let $G = C_p \wr A = K \rtimes A$ where K is the base group of the regular wreath G . Then $K = O_{p',p}(G)$ and $G \in \mathcal{F} = \mathcal{S}_p \mathcal{F}$. Hence $A \simeq G/K = G/O_{p',p}(G) \in F(p)$, a contradiction. Thus $F(p) = \mathcal{F}$.

(2) The inclusion $F(p) \subseteq \mathcal{S}_p \mathcal{H}$ is evident. The inverse inclusion can be proved similarly as the inclusion $\mathcal{F} \subseteq F(p)$ in the proof of (1). \square

We will also use in our proofs the following well-known elementary fact (see for example, [11, Lemma 18.8] or [6, Lemma 3.5.13]).

Lemma 2.10. *If $O_p(G) = 1$ and G has a unique minimal normal subgroup, then there exists a simple $\mathbb{F}_p G$ -module which is faithful for G .*

3. Proofs of the theorems

Proof of Theorem C. (a) Suppose that this assertion is false and let G be a counterexample with minimal order. Let $I = \text{Int}_{\mathcal{F}}(G)$ and $I_1 = \text{Int}_{\mathcal{M}}(G)$. Then $1 < I < G$ and $I_1 \neq G$. Let L be a minimal normal subgroup of G contained in I and $C = C_G(L)$. Then $\pi(L) \subseteq \pi(\mathcal{F})$.

- (1) $IN/N \leq \text{Int}_{\mathcal{F}}(G/N) \leq \text{Int}_{\mathcal{M}}(G/N)$ for any non-identity normal subgroup N of G .

Indeed, by Lemma 2.4(a), we have $IN/N \leq \text{Int}_{\mathcal{F}}(G/N)$. On the other hand, by the choice of G , $\text{Int}_{\mathcal{F}}(G/N) \leq \text{Int}_{\mathcal{M}}(G/N)$.

- (2) $L \not\leq I_1$; in particular, the order of L is divisible by some prime $p \in \pi$.

Suppose that $L \leq I_1$. Then $I_1/L = \text{Int}_{\mathcal{M}}(G/L)$ by Lemma 2.4(e). But by (1), $IL/L = I/L \leq \text{Int}_{\mathcal{F}}(G/L) \leq \text{Int}_{\mathcal{M}}(G/L)$. Hence $I/L \leq I_1/L$ and so $I \leq I_1$, a contradiction. Thus $L \not\leq I_1$. This means that there exists an \mathcal{M} -maximal subgroup M of G such that $L \not\leq M$. Suppose that L is a π' -group.

Then $LM \in \mathcal{G}_\pi \mathcal{M} = \mathcal{M}$, which contradicts the maximality of M . Hence the order of L is divisible by some prime $p \in \pi$.

(3) If $L \leq M < G$, then $L \leq \text{Int}_{\mathcal{M}}(M)$.

By Lemma 2.4(b), $L \leq I \cap M \leq \text{Int}_{\mathcal{F}}(M)$. But since $|M| < |G|$, $\text{Int}_{\mathcal{F}}(M) \leq \text{Int}_{\mathcal{M}}(M)$ by the choice of G . Hence $L \leq \text{Int}_{\mathcal{M}}(M)$.

(4) $G = LU$ for any \mathcal{M} -maximal subgroup U of G not containing L . In particular, $G/L \in \mathcal{M}$.

Indeed, suppose that $LU \neq G$. Then by (3), $L \leq \text{Int}_{\mathcal{M}}(LU)$, which implies that $LU \in \mathcal{M}$ by Lemma 2.4(c). This contradicts the \mathcal{M} -maximality of U . Hence we have (4).

(5) $C_G(L) \cap U = U_G = 1$ for any \mathcal{M} -maximal subgroup U of G not containing L .

Since $C_G(L)$ is normal in G and $G = LU$ by (4), $U_G = C_G(L) \cap U$. Assume that $U_G \neq 1$. Let $U/U_G \leq W/U_G$, where W/U_G is an \mathcal{M} -maximal subgroup of G/U_G . Then by (1), $LU_G/U_G \leq W/U_G$. Hence $G = LU \leq W$ by (4), which means that $G/U_G = W/U_G \in \mathcal{M}$. But by (4), $G/L \in \mathcal{M}$. Therefore $G \simeq G/L \cap U_G \in \mathcal{M}$, and consequently $I = G$, a contradiction. Hence (5) holds.

The final contradiction for (a). Since $L \not\leq I_1$ by (2), there is an \mathcal{M} -maximal subgroup M of G such that $L \not\leq M$. But then $G = LM$ by (4). Since $L \leq I$ and $G \notin \mathcal{F}$, $M \notin \mathcal{F}$ by Lemma 2.4(d). Let H be an \mathcal{F} -critical subgroup of M , V a maximal subgroup of H . We show that $V \in F(p)$. By Lemma 2.4(d), $D = LV \in \mathcal{F}$. Hence $D/O_{p',p}(D) \in F(p)$. First assume that L is a non-abelian group. Then, since p divides $|L|$, $O_{p',p}(D) \cap L = 1$. Hence $O_{p',p}(D) \leq C_G(L)$ and $O_{p',p}(D) \cap V = 1$ by (5). Since \mathcal{F} is hereditary, $F(p)$ is hereditary by Lemma 2.1. Therefore $O_{p',p}(D)V/O_{p',p}(D) \simeq V \in F(p)$. Now assume that L is an abelian p -group. Then $L \leq O_{p',p}(D)$ and $O_{p',p}(D) = L(O_{p',p}(D) \cap V)$. Hence $O_{p',p}(D) \leq M \cap C_G(L) = 1$. It follows that $O_{p',p}(D) = O_p(D)$. Therefore $D/O_p(D) \in F(p) = \mathcal{G}_p F(p)$, which implies that $D \in F(p)$ and so $V \in F(p)$. Therefore H is an $F(p)$ -critical group. Since \mathcal{M} is hereditary and $M \in \mathcal{M}$, $H \in \mathcal{M}$. But then $H \in \mathcal{F}$ since \mathcal{F} satisfies the π -boundary condition in \mathcal{M} by hypothesis. This contradiction completes the proof of (a).

(b) Suppose that every $M(p)$ -critical group G belongs to \mathcal{F} for every $p \in \pi$. First we show that $\mathcal{N} \subseteq \mathcal{M}$. Assume that this is false and let C_q be a group of prime order q with $C_q \notin \mathcal{M}$. Let $p \in \pi$. Then C_q is $M(p)$ -critical and so $C_q \in \mathcal{F} \subseteq \mathcal{M}$ by the hypothesis. This contradiction shows that $\mathcal{N} \subseteq \mathcal{M}$.

Now we show that $\text{Int}_{\mathcal{F}}(G) \leq Z_{\pi\mathcal{M}}(G)$ for every group G . Suppose that this assertion is false and let G be a counterexample with minimal order. Let $I = \text{Int}_{\mathcal{F}}(G)$ and $Z = Z_{\pi\mathcal{M}}(G)$. Then $1 < I < G$ and $Z \neq G$. Let N be a minimal normal subgroup of G and L a minimal normal subgroup of G contained in I . Then $\pi(L) \leq \pi(\mathcal{F})$. We proceed via the following steps.

(1) $IN/N \leq \text{Int}_{\mathcal{F}}(G/N) \leq Z_{\pi\mathcal{M}}(G/N)$.

Indeed, by Lemma 2.4(a), we have $IN/N \leq \text{Int}_{\mathcal{F}}(G/N)$. On the other hand, by the choice of G , $\text{Int}_{\mathcal{F}}(G/N) \leq Z_{\pi\mathcal{M}}(G/N)$.

(2) $L \not\leq Z$; in particular, the order of L is divisible by some prime $p \in \pi$.

Suppose that $L \leq Z$. Then, clearly, $Z/L = Z_{\pi\mathcal{M}}(G/L)$, and $I/L = \text{Int}_{\mathcal{F}}(G/L)$ by Lemma 2.4(e). But by (1), $\text{Int}_{\mathcal{F}}(G/L) \leq Z_{\pi\mathcal{M}}(G/L)$. Hence $I/L \leq Z/L$. Consequently, $I \leq Z$, a contradiction.

(3) If $L \leq M < G$, then $L \leq Z_{\mathcal{M}}(M)$.

By Lemma 2.4(b), $L \leq I \cap M \leq \text{Int}_{\mathcal{F}}(M)$. But since $|M| < |G|$, we have that $\text{Int}_{\mathcal{F}}(M) \leq Z_{\pi\mathcal{M}}(M)$ by the choice of G . Hence $L \leq Z_{\pi\mathcal{M}}(M)$ and so $L \leq Z_{\mathcal{M}}(M)$ since the order of L is divisible by some prime $p \in \pi$ by (2).

(4) $L = N$ is the unique minimal normal subgroup of G .

Suppose that $L \neq N$. Then by (1), $NL/N \leq Z_{\pi\mathcal{M}}(G/N)$. Hence from the G -isomorphism $NL/N \simeq L$ we obtain $L \leq Z$, which contradicts (2).

(5) $L \not\leq \Phi(G)$.

Suppose that $L \leq \Phi(G)$. Then L is a p -group by (2). Let $C = C_G(L)$ and M be any maximal subgroup of G . Then $L \leq M$. Hence $L \leq Z_{\mathcal{M}}(M)$ by (3), so $M/M \cap C \in M(p)$ by Lemmas 2.1(1) and 2.5. If $C \not\leq M$, then $G/C = CM/C \simeq M/M \cap C \in M(p)$. This implies that $L \leq Z$, which contradicts (2). Hence $C \leq M$ for all maximal subgroups M of G . It follows that C is nilpotent. Then in view of (4), C is a p -group since C is normal in G . Hence for every maximal subgroup M of G we have $M \in \mathcal{G}_p M(p) = M(p)$. But since $M(p) \subseteq \mathcal{M}$, $G \notin M(p)$ (otherwise $G \in \mathcal{M}$ and so $G = Z$). This shows that G is an $M(p)$ -critical group. Therefore $G \in \mathcal{F}$ by the hypothesis. But since $\mathcal{F} \subseteq \mathcal{M}$, we have $G \in \mathcal{M}$ and so $G = Z$, a contradiction. Thus (5) holds.

(6) $C = C_G(L) \leq L$ (this follows from (4), (5) and Theorem 15.6 in [4, Chapter A]).

(7) If $L \leq M < G$, then $M \in M(p)$.

First by (3), $L \leq Z_{\mathcal{M}}(M)$. If $L = C$, then $M/L = M/M \cap C \in M(p)$ by Lemma 2.5, which implies that $M \in \mathcal{G}_p M(p) = M(p)$ since L is a p -group by (2). Now suppose that L is a non-abelian group. Let $1 = L_0 < L_1 < \dots < L_n = L$ be a chief series of M below L . Let $C_i = C_M(L_i/L_{i-1})$ and $C_0 = C_1 \cap \dots \cap C_n$. Since $L \leq Z_{\mathcal{M}}(M)$, $M/C_i \in M(p)$ for all $i = 1, \dots, n$. It follows that $M/C_0 \in M(p)$. Since $C = 1$ by (4) and (6), for any minimal normal subgroup R of M we have $R \leq L$. Suppose that $C_0 \neq 1$ and let R be a minimal normal subgroup of M contained in C_0 . Then $R \leq L$ and $R \leq C_M(H/K)$ for each chief factor H/K of M . Thus $R \leq F(M)$ is abelian and hence L is abelian. This contradiction shows that $C_0 = 1$. Consequently, $M \in M(p)$.

(8) There exists a subgroup U of G such that $U \in \mathcal{F}$ and $LU = G$.

Indeed, suppose that every maximal subgroup of G not containing L belongs to $M(p)$. Then by (7), G is an $M(p)$ -critical group. Hence $G \in \mathcal{F}$ by the hypothesis. But then $I = G$, a contradiction. Hence there exists a maximal subgroup M of G such that $G = LM$ and $M \notin M(p)$. Take an $M(p)$ -critical subgroup U of M . Then in view of (7), $LU = G$ and $U \in \mathcal{F}$ by the hypothesis.

(9) The final contradiction for (b).

Since $L \leq I$ and $G/L = UL/L \simeq U/U \cap L \in \mathcal{F}$ by (8), it follows from Lemma 2.4(c) that $G \in \mathcal{F}$ and so $G = I$. The final contradiction shows that $\text{Int}_{\mathcal{F}}(G) \leq Z_{\pi\mathcal{M}}(G)$ for every group G . The second assertion of (b) can be proved similarly. The theorem is proved. \square

Proofs of Theorems A and B. Since $Z_{\mathcal{F}}(G) \leq \text{Int}_{\mathcal{F}}(G)$ by Lemma 2.4(g), the sufficiency is a special case, when $\mathcal{F} = \mathcal{M}$, of Theorem C(b). Now suppose that the equality $Z_{\pi\mathcal{F}}(G) = \text{Int}_{\mathcal{F}}(G)$ holds for each (soluble) group G .

First we show that $\mathcal{N} \subseteq \mathcal{F}$. Let F be the canonical local satellite of \mathcal{F} . Suppose that for some group C_q of prime order q we have $C_q \notin \mathcal{F}$. Let $p \in \pi$ and $G = PC_q$, where P is a simple $\mathbb{F}_p C_q$ -module P which is faithful for C_q . Then $P = \text{Int}_{\mathcal{F}}(G)$ and $Z_{\mathcal{F}}(G) = 1$ since $F(p) \subseteq \mathcal{F}$. This contradiction shows that $\mathcal{N} \subseteq \mathcal{F}$.

Now we show that $\mathcal{G}_{\pi'}\mathcal{F} = \mathcal{F}$ ($\mathcal{S}_{\pi'}\mathcal{F} = \mathcal{F}$, respectively). The inclusion $\mathcal{F} \subseteq \mathcal{G}_{\pi'}\mathcal{F}$ is evident. Suppose that $\mathcal{G}_{\pi'}\mathcal{F} \not\subseteq \mathcal{F}$ ($\mathcal{S}_{\pi'}\mathcal{F} \not\subseteq \mathcal{F}$) and let G be a group of minimal order in $\mathcal{G}_{\pi'}\mathcal{F} \setminus \mathcal{F}$ (in $\mathcal{S}_{\pi'}\mathcal{F} \setminus \mathcal{F}$, respectively). Then $G^{\mathcal{F}}$ is the unique minimal normal subgroup of G and $G^{\mathcal{F}}$ is a π' -group. Hence

$$G^{\mathcal{F}} \leq Z_{\pi\mathcal{F}}(G) = \text{Int}_{\mathcal{F}}(G).$$

It follows from Lemma 2.4(c) that $G \in \mathcal{F}$. This contradiction shows that $\mathcal{G}_{\pi'}\mathcal{F} = \mathcal{F}$ ($\mathcal{S}_{\pi'}\mathcal{F} = \mathcal{F}$, respectively).

Finally, we show that \mathcal{F} satisfies the π -boundary condition (the π -boundary condition in the class \mathcal{S} , respectively). Suppose that this is false. Then for some $p \in \pi$, the set of all (soluble) $F(p)$ -critical groups A with $A \notin \mathcal{F}$ is non-empty. Let us choose in this set a group A with minimal $|A|$. Then by Lemma 2.6, $A^{\mathcal{F}}$ is the unique minimal normal subgroup of G and $O_p(A) = 1 = \Phi(A)$. Hence by Lemma 2.10, there exists a simple $\mathbb{F}_p A$ -module P which is faithful for A . Let $G = P \rtimes A$ and M be any maximal subgroup of G . If $P \not\leq M$, then $M \simeq G/P \simeq A \notin \mathcal{F}$. On the other hand, if $P \leq M$, then $M = M \cap PA = P(M \cap A)$, where $M \cap A$ is a maximal subgroup of A . Hence $M \cap A \in F(p)$ and so $M \in \mathcal{G}_p F(p) = F(p) \subseteq \mathcal{F}$. Therefore P is contained in the intersection of all \mathcal{F} -maximal subgroups of G . Then $P \leq Z_{\pi\mathcal{F}}(G)$ by our assumption on \mathcal{F} . It follows that $A \simeq G/P = G/C_G(P) \in F(p) \subseteq \mathcal{F}$ by Lemma 2.2(1) and Lemma 2.5. But this contradicts the choice of A . Therefore \mathcal{F} satisfies the π -boundary condition (\mathcal{F} satisfies the π -boundary condition in the class \mathcal{S}). The theorems are proved. \square

In view of Theorems A, B and C we have

Corollary 3.1. Let \mathcal{F} be a hereditary saturated formation with $(1) \neq \mathcal{F} \neq \mathcal{S}$. Then the equality $\text{Int}_{\mathcal{F}}(G) = Z_{\mathcal{F}}(G)$ holds for each group G if and only if $\mathcal{N} \subseteq \mathcal{F}$ and \mathcal{F} satisfies the boundary condition.

Corollary 3.2. Let \mathcal{F} be a hereditary saturated formation of soluble groups with $(1) \neq \mathcal{F} \neq \mathcal{S}$. Then the equality $\text{Int}_{\mathcal{F}}(G) = Z_{\mathcal{F}}(G)$ holds for each soluble group G if and only if $\mathcal{N} \subseteq \mathcal{F}$ and \mathcal{F} satisfies the boundary condition in the class \mathcal{S} .

Note that Corollary 3.2 also follows directly from [3, Main Theorem].

Proof of Theorem D. We will prove the theorem by induction on $|G|$. If $G \in \mathcal{F}$, then

$$\text{Int}_{\mathcal{F}}^*(G) = G = \text{Int}_{\mathcal{F}}(G).$$

We may, therefore, assume that $G \notin \mathcal{F}$. Let $I = \text{Int}_{\mathcal{F}}(G)$, $I^* = \text{Int}_{\mathcal{F}}^*(G)$ and N be a minimal normal subgroup of G . Then $I \leq I^*$. Hence we may assume that $I^* \neq 1$.

(1) $I^*N/N \leq \text{Int}_{\mathcal{F}}^*(G/N)$.

If U/N is a non- K - \mathcal{F} -subnormal \mathcal{F} -maximal subgroup of G/N , then for some non- K - \mathcal{F} -subnormal \mathcal{F} -maximal subgroup U_0 of G we have $U = U_0N$ by Lemma 2.8(i). Let

$$\text{Int}_{\mathcal{F}}^*(G/N) = U_1/N \cap \cdots \cap U_t/N,$$

where U_i/N is a non- K - \mathcal{F} -subnormal \mathcal{F} -maximal subgroup of G/N for all $i = 1, \dots, t$. Let V_i be a non- K - \mathcal{F} -subnormal \mathcal{F} -maximal subgroup of G such that $U_i = V_iN$. Then $I^* \leq V_1 \cap \cdots \cap V_t$. Hence $I^*N/N \leq \text{Int}_{\mathcal{F}}^*(G/N)$.

(2) If $N \leq I^*$, then $\text{Int}_{\mathcal{F}}^*(G/N) = I^*/N$.

By Lemma 2.8(i), it is enough to prove that if U is a non- K - \mathcal{F} -subnormal \mathcal{F} -maximal subgroup of G , then U/N is a non- K - \mathcal{F} -subnormal \mathcal{F} -maximal subgroup of G/N . Let $U/N \leq X/N$, where X/N is a non- K - \mathcal{F} -subnormal \mathcal{F} -maximal subgroup of G/N . By Lemma 2.8(i), $X = U_0N$ for some non- K - \mathcal{F} -subnormal \mathcal{F} -maximal subgroup U_0 of G . But since $N \leq U_0$, $U/N \leq U_0/N$ and so $U = U_0$. Thus $U/N = X/N$.

(3) $I^* \cap H \leq \text{Int}_{\mathcal{F}}^*(H)$ for any subgroup H of G .

Let V be an arbitrary non- K - \mathcal{F} -subnormal \mathcal{F} -maximal subgroup of H . Then $V = H \cap U$ for some non- K - \mathcal{F} -subnormal \mathcal{F} -maximal subgroup U of G by Lemma 2.8(ii). Thus there are non- K - \mathcal{F} -subnormal \mathcal{F} -maximal subgroups U_1, \dots, U_t of G such that

$$\text{Int}_{\mathcal{F}}^*(H) = U_1 \cap \cdots \cap U_t \cap H.$$

This induces that $I^* \cap H \leq \text{Int}_{\mathcal{F}}^*(H)$.

(4) If $E = DV$ for some normal subgroup D of G contained in I^* and some K - \mathcal{F} -subnormal subgroup $V \in \mathcal{F}$ of G , then $E \in \mathcal{F}$.

First note that $R \leq \text{Int}_{\mathcal{F}}^*(E)$ by (3). On the other hand, by Lemma 2.7(i), V is a K - \mathcal{F} -subnormal subgroup of E . Hence we need only consider the case when $G = E$. Assume that $G \notin \mathcal{F}$. Let R be any minimal normal subgroup of G . Then $(DR/R)(VR/R) = G/R$, where $DR/R \leq \text{Int}_{\mathcal{F}}^*(G/R)$ by (1), and $VR/R \simeq V/V \cap R \in \mathcal{F}$ is a K - \mathcal{F} -subnormal subgroup of G/R . Hence by induction we have $G/R \in \mathcal{F}$. This implies that R is the only minimal normal subgroup of G and so $R = G^{\mathcal{F}} \leq I^*$. Let W be a minimal supplement of R in G . Then $W \in \mathcal{F}$ by Lemma 2.3(i). Let $W \leq M$, where M is an \mathcal{F} -maximal subgroup of G . If M is not K - \mathcal{F} -subnormal in G , then $R \leq M$. Thus $G = RW = RM = M \in \mathcal{F}$, a contradiction. This shows that M is K - \mathcal{F} -subnormal in G . But then there is a proper subgroup X of G such that $M \leq X$ and either X is normal in G or $R = G^{\mathcal{F}} \leq X$. In both of these cases, we have that $G = RM = RX = X < G$, a contradiction. Hence we have (4).

Conclusion. Let R be a minimal normal subgroup of G contained in I^* . If $R \leq I$, then $I/R = \text{Int}_{\mathcal{F}}(G/R)$ by Lemma 2.4(e), and $I^*/R = \text{Int}_{\mathcal{F}}^*(G/R)$ by (2). Therefore by induction, $\text{Int}_{\mathcal{F}}^*(G/R) = \text{Int}_{\mathcal{F}}(G/R)$. It follows that $I = I^*$.

Finally, suppose that $R \not\leq I$. Then $R \not\leq U$ for some \mathcal{F} -maximal subgroup U of G . Let $E = RU$. Then $R \leq \text{Int}_{\mathcal{F}}^*(E)$ by (3). On the other hand, it is clear that U is a K - \mathcal{F} -subnormal subgroup of G . Hence by (4), $E \in \mathcal{F}$. But then $E = U$, which implies $R \leq U$, a contradiction. The theorem is proved. \square

4. Applications and remarks

Applications of Theorems A, B and D. We say that \mathcal{F} satisfies the p -boundary condition if \mathcal{F} satisfies the $\{p\}$ -boundary condition in the class of all groups.

Lemma 4.1. Let $\mathcal{F} = LF(F)$, where F is the canonical local satellite of \mathcal{F} . Suppose that for some prime p we have $F(p) = \mathcal{F}$. Then \mathcal{F} does not satisfy the p -boundary condition.

Proof. Indeed, in this case every \mathcal{F} -critical group is also $F(p)$ -critical. \square

A group G is called π -closed if G has a normal Hall π -subgroup.

Proposition 4.2. The formation \mathcal{F} of all π -closed groups satisfies the π' -boundary condition, but \mathcal{F} does not satisfy the p -boundary condition for any $p \in \pi$.

Proof. Let $\mathcal{F} = \mathcal{G}_\pi \mathcal{G}_{\pi'}$ be the formation of all π -closed groups, F the canonical local satellite of \mathcal{F} . Then $F(p) = \mathcal{F}$ for all $p \in \pi$, and $F(p) = \mathcal{G}_{\pi'}$ for all primes $p \in \pi'$ by Theorem 3.1.20 in [6]. Hence \mathcal{F} satisfies the π' -boundary condition and does not satisfy the p -boundary condition for any $p \in \pi$ by Lemma 4.1. \square

Lemma 4.3. Let $\{\mathcal{F}_i \mid i \in I\}$ be any set of non-empty saturated formations and $\mathcal{F} = \bigcap_{i \in I} \mathcal{F}_i$.

(1) If for each $i \in I$, \mathcal{F}_i satisfies the p -boundary condition, then \mathcal{F} satisfies the p -boundary condition.

(2) Suppose that $I = \{1, 2\}$, F_i is the canonical local satellite of \mathcal{F}_i and that there is a set π of primes satisfying the following conditions:

(a) \mathcal{F}_1 satisfies the π -boundary condition, and for any $p \in \pi$, we have $F_1(p) \subseteq \mathcal{F}_2 = F_2(p)$ and every $F_1(p)$ -critical group belongs to \mathcal{F}_2 .

(b) \mathcal{F}_2 satisfies the π' -boundary condition, and for any $p \in \pi'$, we have $F_2(p) \subseteq \mathcal{F}_1 = F_1(p)$ and every $F_2(p)$ -critical group belongs to \mathcal{F}_1 .

Then \mathcal{F} satisfies the boundary condition.

Proof. (1) Let F_i be the canonical local satellite of \mathcal{F}_i and F the canonical local satellite of \mathcal{F} . If $f(p) = \bigcap_{i \in I} F_i(p)$, then $F(p) = \mathcal{G}_p f(p)$ by Theorem 3.3 in [10, Chapter 1]. Now let G be any $F(p)$ -critical group, $i \in I$. Since $F(p) \subseteq F_i(p)$, all maximal subgroup of G belongs to $F_i(p)$. Hence $G \in \mathcal{F}_i$ since $F_i(p) \subseteq \mathcal{F}_i$ and \mathcal{F}_i satisfies the p -boundary condition. This implies that $G \in \mathcal{F}$ and therefore \mathcal{F} satisfies the p -boundary condition.

(2) In this case, $F(p) = F_1(p)$ for all $p \in \pi$ and $F(p) = F_2(p)$ for all $p \in \pi'$, where F is the canonical local satellite of \mathcal{F} . Hence if $p \in \pi$ and G is an $F(p)$ -critical group, then $G \in \mathcal{F}$ by hypothesis (a). This shows that \mathcal{F} satisfies the π -boundary condition. Similarly we see that \mathcal{F} satisfies the π' -boundary condition. \square

A group G is called p -decomposable if there exists a subgroup H of G such that $G = P \times H$ for some (and hence the unique) Sylow p -subgroup P of G .

Corollary 4.4. The formation of all p -decomposable groups satisfies the boundary condition.

Proof. Let \mathcal{F} be the formation of all p -decomposable groups. Then $\mathcal{F} = \mathcal{G}_{p'} \mathcal{G}_p \cap \mathcal{G}_p \mathcal{G}_{p'}$. Hence the assertion follows from Proposition 4.2 and Lemma 4.3(2). \square

From Corollary 4.4 and Theorem A we get

Corollary 4.5. Let D be the intersection of all maximal p -decomposable subgroups of G . Then D is the largest normal subgroup of G satisfying $D = O_{p'}(D) \times O_p(D)$, and G induces the trivial automorphisms group on every chief factor of G below $O_p(D)$ and a π' -group of automorphisms on every chief factor of G below $O_{p'}(D)$.

Since a p -nilpotent group is p' -closed, the following result directly follows from Proposition 4.2.

Corollary 4.6. The formation of all p -nilpotent groups satisfies the p -boundary condition.

From Corollary 4.6 and Theorem A, we have

Corollary 4.7. *Let D be the intersection of all maximal p -nilpotent subgroups of a group G . Then D is the largest normal subgroup of G satisfying $O_{p'}(D) = O_{p'}(G)$, and $D/O_{p'}(G) \leq Z_\infty(G/O_{p'}(G))$.*

Note that another proof of Corollary 4.7 was obtained in the paper [3].

Remark 4.8. If $\pi \neq \{2\}$, then the formation \mathcal{F} of all π -supersoluble groups does not satisfy the π -boundary condition in the class of all soluble groups. Indeed, let F be the canonical local satellite of \mathcal{F} . Then $F(p) = \mathcal{N}_p\mathcal{A}(p-1)$, where $\mathcal{A}(p-1)$ is the formation of all abelian groups of exponent dividing $p-1$ [4, p. 358], for all $p \in \pi$ and $F(p) = \mathcal{F}$ for all primes $q \notin \pi$ (see Example 3.4(e) and Theorem 4.8 in [4, Chapter IV]). Let $2 \neq p \in \pi$ and q be any prime with q divides $p-1$. Let $G = Q \rtimes C$, where C is a group of order p and Q is an $\mathbb{F}_q C$ -module which is faithful for C . Then G is soluble, but G is a non-supersoluble $F(p)$ -critical groups.

Proposition 4.9. *Let $\mathcal{F} = \mathcal{NL}$.*

(i) *If \mathcal{L} is a hereditary saturated formation satisfying the boundary condition in the class of all soluble groups, then \mathcal{F} satisfies the boundary condition in the class of all soluble groups.*

(ii) *If \mathcal{L} is a formation of nilpotent groups with $\pi(\mathcal{L}) = \mathbb{P}$, then \mathcal{F} satisfies the boundary condition.*

Proof. Let F be the canonical local satellite of \mathcal{F} . Then by Lemma 2.9(2), $F(p) = \mathcal{G}_p\mathcal{L}$ for all primes p . Assume that \mathcal{F} does not satisfy the boundary condition (\mathcal{F} does not satisfy the boundary condition in the class of all soluble groups). Then for some prime p , the set of all $F(p)$ -critical (soluble) groups A with $A \notin \mathcal{F}$ is non-empty. Let G be a group of minimal order in this set. Then $L = G^\mathcal{F}$ is the unique minimal normal subgroup of G and $O_p(G) = 1 = \Phi(G)$ by Lemma 2.6.

First suppose that G is soluble. Then $L = C_G(L)$ is a q -group for some prime $q \neq p$ and $G = L \rtimes M$ for some maximal subgroup M of G with $O_q(M) = 1$ by [4, Chapter A, Theorem 15.6]. Let M_1 be any maximal subgroup of M . Then $LM_1 \in F(p)$, and so $LM_1 \in \mathcal{L}$ since $L = C_G(L)$ is a q -group.

Suppose that \mathcal{L} satisfies the condition of (i) and let L be the canonical local satellite of \mathcal{L} . Since $LM_1 \in \mathcal{L}$,

$$LM_1/O_{q',q}(LM_1) = LM_1/O_q(LM_1) = LM_1/LO_q(M_1) \simeq M_1/M_1 \cap LO_q(M_1) \in L(q) = \mathcal{G}_p L(q).$$

Hence every maximal subgroup of M belongs to $L(q)$. Since \mathcal{L} satisfies the boundary condition in the class of all soluble groups, we obtain that $M \in \mathcal{L}$ and consequently $G \in \mathcal{F}$. This contradiction completes the proof of (i).

Suppose that \mathcal{L} satisfies the condition (ii). Let M_1 be a normal maximal subgroup of M . Since $LM_1 \in \mathcal{L}$ and $L = C_G(L)$, we have $M_1 = 1$. This implies that $|M|$ is prime. Hence $M \in \mathcal{L}$ since $\pi(\mathcal{L}) = \mathbb{P}$. But then $G \in \mathcal{F} = \mathcal{NL}$. Therefore G is not soluble.

Let $q \neq p$ be any prime divisor of $|G|$. Suppose that G is not q -nilpotent. Then G has a q -closed \mathcal{N} -critical subgroup $H = Q \rtimes R$ by [7, IV, Theorem 5.4], where Q is a Sylow q -subgroup of H , R is a cyclic Sylow r -subgroup of H . Since G is not soluble, $H \neq G$. Hence $H \leq M \in F(p) = \mathcal{G}_p\mathcal{L}$ for some maximal subgroup M of G . Since $M \in \mathcal{G}_p\mathcal{L}$, $M^\mathcal{L} \leq O_p(M)$ and hence $H^\mathcal{L} \leq Q \cap O_p(H) = 1$. This shows that H is nilpotent. This contradiction shows that G is q -nilpotent for all primes $q \neq p$. This induces that $G^\mathcal{N}$ is a p -subgroup of G and thereby G is soluble. This contradiction completes the proof of (ii). \square

Remark 4.10. The condition “ \mathcal{L} is a hereditary saturated formation satisfying the boundary condition in the class of all soluble groups” cannot be omitted in the statement (i). Indeed, let $\mathcal{F} = \mathcal{NU}$ and $G = P \rtimes A_4$, where P is a simple $\mathbb{F}_3 A_4$ -module P which is faithful for A_4 . Let F be the canonical local satellite of \mathcal{F} . Then $F(2) = \mathcal{G}_2\mathcal{U}$ by Lemma 2.9(2). Therefore G is an $F(2)$ -critical group and $G \notin \mathcal{F}$. Thus G does not satisfy the boundary condition in the class of all soluble groups.

Corollary 4.11. *Let \mathcal{F} be the class of all groups with $G' \leq F(G)$. Then \mathcal{F} satisfies the boundary condition.*

From Corollary 4.11 and Theorem A, we obtain

Corollary 4.12. *Let D be the intersection of all maximal subgroups H of G with the property $H' \leq F(H)$. Then D is the largest normal subgroup of G such that $D' \leq F(D)$ and G induces an abelian group of automorphisms on every chief factor of G below D .*

Note that Corollary 4.12 can be also found in [3, Corollary 7 and Remark 4].

Following [4, Chapter VII, Definition 6.9] we write $l(G)$ to denote the nilpotent length of the group G . Recall that \mathcal{N}^r is the product of r copies of \mathcal{N} ; \mathcal{N}^0 is the class of groups of order 1 by definition. It is well known that \mathcal{N}^r is the class of all soluble groups G with $l(G) \leq r$. It is also known that \mathcal{N}^r is a hereditary saturated formation (see, for example, [4, p. 358]). Hence from Proposition 4.9 we get

Corollary 4.13. *Let $\mathcal{F} = \mathcal{N}^r \mathcal{L}$ ($r \in \mathbb{N}$), where \mathcal{L} is a subformation of the formation of all abelian groups with $\pi(\mathcal{L}) = \mathbb{P}$. Then \mathcal{F} satisfies the boundary condition in the class of all soluble groups.*

From Proposition 4.9 and Theorem A we get

Corollary 4.14. *Let D be the intersection of all maximal metanilpotent subgroups of G . Then D is the largest normal subgroup of G such that D is metanilpotent and G induces a nilpotent group of automorphisms on every chief factor of G below D .*

It is clear that every subnormal subgroup is a K - \mathcal{F} -subnormal subgroup as well. On the other hand, in the case when $\mathcal{N} \subseteq \mathcal{F}$, every K - \mathcal{F} -subnormal subgroup of any soluble subgroup G is \mathcal{F} -subnormal in G . Hence from Theorem D and the above corollaries we get

Corollary 4.15. *Let \mathcal{F} be the class of all groups G with $G' \leq F(G)$. Then:*

- (i) *The subgroup $Z_{\mathcal{F}}(G)$ may be characterized as the intersection of all non-subnormal \mathcal{F} -maximal subgroups of G , for each group G .*
- (ii) *The subgroup $Z_{\mathcal{F}}(G)$ may be characterized as the intersection of all non- \mathcal{F} -subnormal \mathcal{F} -maximal subgroups of G , for each soluble group G .*

Corollary 4.16. *Let \mathcal{F} be one of the following formations:*

- (1) *the class of all nilpotent groups (Baer [1]);*
 - (2) *the class of all groups G with $G' \leq F(G)$ (Skiba [13]);*
 - (3) *the class of all p -decomposable groups (Skiba [12]).*
- Then for each group G , the equality $\text{Int}_{\mathcal{F}}(G) = Z_{\mathcal{F}}(G)$ holds.*

Corollary 4.17. (Sidorov [9].) *Let \mathcal{F} be the class of all soluble groups G with $l(G) \leq r$ ($r \in \mathbb{N}$). Then for each soluble group G , the equality $Z_{\mathcal{F}}(G) = \text{Int}_{\mathcal{F}}(G)$ holds.*

Let $p_1 > p_2 > \dots > p_r$ be the distinct primes dividing $|G|$, P_i a Sylow p_i -subgroup of G . Then G is said to satisfy the Sylow tower property if all subgroups $P_1, P_1 P_2, \dots, P_1 P_2 \dots P_{r-1}$ are normal in G . It is well known that every supersoluble group satisfies the Sylow tower property. As an application of Theorem C(a) we have the following result.

Proposition 4.18. *Let \mathcal{F} be the formation of all supersoluble groups and \mathcal{M} be the formation of all groups satisfying the Sylow tower property. Then $\text{Int}_{\mathcal{F}}(G) \leq \text{Int}_{\mathcal{M}}(G)$ for any group G .*

Proof. Let F be the canonical local satellite of \mathcal{F} . Then $F(p) = \mathcal{G}_p \mathcal{A}(p-1)$ for primes p (see Remark 4.8). Let G be any $F(p)$ -critical group satisfying the Sylow tower property. We show that $G \in \mathcal{F}$. Let q be the largest prime dividing $|G|$, P the Sylow q -subgroup of G . If $G = P$, then clearly $G \in \mathcal{F}$. Let $P \neq G$. Then every Sylow subgroup of G belongs to $F(p)$. Hence $q = p$ and if E is a Hall p' -subgroup of G , then $E \in \mathcal{A}(p-1)$. But then $G \in F(p)$, a contradiction. Therefore $G \in \mathcal{F}$. This shows that \mathcal{F} satisfies the boundary condition in \mathcal{M} . Hence, in view of Theorem C(a), we have $\text{Int}_{\mathcal{F}}(G) \leq \text{Int}_{\mathcal{M}}(G)$. \square

Remark 4.19. If $\mathcal{F} \subseteq \mathcal{M}$ are hereditary saturated formations, then for every group G we have $Z_{\mathcal{F}}(G) \leq Z_{\mathcal{M}}(G)$ and $Z_{\mathcal{F}}(G) \leq \text{Int}_{\mathcal{F}}(G)$ by Lemma 2.4(g). Therefore, if \mathcal{F} satisfies the boundary condition, then $\text{Int}_{\mathcal{F}}(G) \leq \text{Int}_{\mathcal{M}}(G)$ for every group G . But, by Remark 4.8, \mathcal{F} does not necessarily satisfy the boundary condition. Hence we cannot deduce Proposition 4.18 from Theorem A.

Lemma 4.20. Let $\mathcal{F} = LF(F)$ be a saturated formation, where F is the canonical local satellites of \mathcal{F} . Let $\mathcal{M} = \mathcal{F} \cap \mathcal{S}$ and M be the canonical local satellites of \mathcal{M} . Then $M(p) = F(p) \cap \mathcal{S}$ for all primes p .

Proof. It is clear that $\mathcal{M} = LF(F_1)$, where $F_1(p) = F(p) \cap \mathcal{S}$ for all primes p . On the other hand, for any prime p we have $F_1(p) \subseteq \mathcal{M}$, and $F_1(p) = \mathcal{G}_p F_1(p)$ since

$$\mathcal{G}_p F_1(p) = \mathcal{G}_p (F(p) \cap \mathcal{S}) \subseteq F(p) \cap \mathcal{S}.$$

Hence $F_1 = M$ is the canonical local satellite of \mathcal{M} . \square

Lemma 4.21. Let $(1) \neq \mathcal{F} = LF(F)$ be a saturated formation and $\mathcal{M} = \mathcal{F} \cap \mathcal{S}$. Let $\pi \subseteq \pi(\mathcal{F})$. Then \mathcal{F} satisfies the boundary condition in the class \mathcal{S} of all soluble groups if and only if \mathcal{M} satisfies the boundary condition in \mathcal{S} .

Proof. Let F and M be the canonical local satellites of the formations \mathcal{F} and \mathcal{M} , respectively. Let $p \in \pi$. Then $M(p) = F(p) \cap \mathcal{S}$ by Lemma 4.20. Hence a soluble group G is $F(p)$ -critical group if and only if G is $M(p)$ -critical. On the other hand, the soluble group G belongs to \mathcal{F} if and only if G belongs to \mathcal{M} . \square

In view of Lemmas 4.20 and 4.21, we may similarly prove the following general form of Theorem B.

Theorem 4.22. Let \mathcal{F} be a saturated formation such that $\mathcal{M} = \mathcal{F} \cap \mathcal{S}$ is hereditary and $(1) \neq \mathcal{M} \neq \mathcal{S}$. Let $\pi \subseteq \pi(\mathcal{F})$. Then the equality

$$Z_{\pi \mathcal{F}}(G) = \text{Int}_{\mathcal{F}}(G)$$

holds for each soluble group G if and only if $\mathcal{N} \subseteq \mathcal{F} = \mathcal{S}_{\pi'} \mathcal{F}$ and \mathcal{F} satisfies the π -boundary condition in the class of all soluble groups.

Acknowledgment

We cordially thank the referee for his/her careful reading and very helpful suggestions. In fact, some results and proofs were modified by the referee, and the ideas of Proposition 4.9 and Corollary 4.14 belong to the referee.

References

- [1] R. Baer, Group elements of prime power index, Trans. Amer. Math. Soc. 75 (1953) 20–47.
- [2] A. Ballester-Bolinches, L.M. Ezquerro, Classes of Finite Groups, Springer, Dordrecht, 2006.
- [3] J.C. Beidleman, H. Heineken, A note on intersections of maximal \mathcal{F} -subgroups, J. Algebra 333 (2010) 120–127.
- [4] K. Doerk, T. Hawkes, Finite Soluble Groups, Walter de Gruyter, Berlin, New York, 1992.

- [5] D. Gorenstein, *Finite Groups*, Harper & Row Publishers, New York, Evanston, London, 1968.
- [6] W. Guo, *The Theory of Classes of Groups*, Science Press/Kluwer Academic Publishers, Beijing, New York, Dordrecht, Boston, London, 2000.
- [7] B. Huppert, *Endliche Gruppen I*, Springer-Verlag, Berlin, Heidelberg, New York, 1967.
- [8] O.H. Kegel, Untergruppenverbände endlicher Gruppen, die den subnormalteilerverband each enthalten, *Arch. Math. (Basel)* 30 (3) (1978) 225–228.
- [9] A.V. Sidorov, On properties of the \mathcal{F} -hypercentre of finite groups, *Probl. in Algebra* 10 (1996) 141–143.
- [10] L.A. Shemetkov, *Formations of Finite Groups*, Nauka, Moscow, 1978.
- [11] L.A. Shemetkov, A.N. Skiba, *Formations of Algebraic Systems*, Nauka, Moscow, 1989.
- [12] A.N. Skiba, On the intersection of all maximal \mathcal{F} -subgroups of a finite group, *J. Algebra* 343 (2011) 173–182.
- [13] A.N. Skiba, On the \mathcal{F} -hypercentre and the intersection of all \mathcal{F} -maximal subgroups of a finite group, *J. Pure Appl. Algebra* 216 (4) (2012) 789–799, <http://dx.doi.org/10.1016/j.jpaa.2011.10.006>.